KNOWN AND PROBABLE KARST IN OHIO
Revised 2006

Drift-thickness and glacial boundary information are added to the map to depict the relationship between drift thickness and active karst processes, and the increasing density of karst features beyond the glacial margins.



KNOWN AND PROBABLE KARST IN OHIO

By

Richard R. Pavey, Dennis N. Hull, C. Scott Brockman, Gregory A. Schumacher, David A. Stith, E. Mac Swinford, Donovan M. Powers, Terry L. Sole, Kim E. VorBau, Kevin D. Kulhal, Emily E. Evans, Ernie R. Shelley, and Robert G. Van Horn

Karst terrains exist in many Ohio counties where dissolution processes in carbonate rocks (limestone and dolomite) and gypsum (calcium sulfate) have produced networks of subsurface conduits. The rapid transmittal of pollutants in the environment through karst is a major concern. In Ohio, karst areas are the most vulnerable natural settings for ground-water contamination. Other karst related problems include water delays and additional expenses in construction projects, and the potential for sinkhole collapse in areas of urbanized bedrock. Although it has long been known that Ohio has cave and sinkhole areas, there had been no comprehensive mapping of the state's karst terrains before this investigation.

In 1995, in anticipation of the statewide search for a low-level radioactive-waste disposal site, the Ohio General Assembly enacted site-screening criteria. These criteria included Section 3747.12 of the Ohio Revised Code, which states, "The disposal site shall not be located in areas of known or probable karst. As used in Division (M)(3) of this Section, 'karst' means a terrain with an assemblage of landforms such as sinkholes and caves that are the result of dissolution of carbonate bedrock or solution of gypsum." To meet this criterion, the Ohio Division of Geological Survey, through a grant from the Midwest Compact administered by the Ohio Low-Level Radioactive Waste Disposal Facility Development Authority, conducted an investigation in 1997 to document vertically expressed karst features in the state.

The investigation began with a review of cave inventories, theses, and scientific literature for field-documented karst locations. These locations were defined as known karst areas. Next, the locations of known karst features were digitized. The boundaries of known karst features was the Ohio Cave Inventory provided by Dr. Horton H. Hobbs II of Wittenberg University. Next, the locations of additional karst features were identified through examination of geomorphic features on 7.5-minute topographic maps in carbonate-rock terrains, analysis of county soil-survey reports, and interpretation of aerial photographs. Areas where karst features are likely to occur were defined as areas of the south terrain, but for clarity are not shown on this map. Field inspection of many indicated karst features documented more known karst locations and verified the mapping methods.

Probable karst areas were defined as areas that: (1) lie within a half mile of a known or indicated karst location, and (2) are underlain by carbonate or gypsiferous bedrock with overburden generally of less than 200 feet of non-carbonate bedrock and/or unconsolidated material as shown by comparison of 7.5-minute bedrock-topography and bedrock-geology maps to surface topography. Known karst locations and probable karst areas on the south terrain were digitized and are shown on this 1:500,000-scale map.

The May 2006 revision of this map is the first to portray information on the thickness of glacial drift and alluvium in areas of karst in Ohio. The drift-thickness map (figure 1, fig. 2) illustrates that karst terrains in Ohio is generally limited to areas of carbonate rock (limestone and dolomite) overlain by less than 200 feet of glacial drift and/or alluvium, and 2) isolated areas of the state where drift-thickness/bedrock-lithology conditions are most conducive to karst development, even though reconnaissance mapping may not have identified karst features. Areas where drift is thicker than 200 feet and areas where bedrock crops out are depicted on this map occur where bedrock is indicated to be carbonate by more than 200 feet of drift and/or by outside the boundaries of bedrock lithologies most susceptible to karst development. Such separate outliers from the bedrock used to map karst terrains are the Devonian age Ohio Shale and Silurian age Salina Group. Karst features were not available in 1997 when probability areas were originally determined, and 3) that have occasionally developed in areas where soluble bedrock underlies as much as 20 or more feet of relatively insoluble bedrock (e.g., noncalcareous shale).

Although Ohio has not been regarded as having a significant amount of karst, this investigation revealed many areas of the state that contain well-developed karst terrains, which occur mostly in four geographic regions of the state: (1) north-central Ohio, where a broad band of solution-collapse depressions extends from northeastern Seneca County to the Ohio River in Scioto County; (2) northeastern Ohio, in parts of Ashtabula, Lake, Geauga, Summit, and Portage counties; (3) southern Ohio, in carbonate outcrops lying beyond the Wisconsinan glacial boundary; and (4) west-central Ohio, in the Bellefontaine upland of Logan and Champaign Counties.

Several relationships between geologic units and the distribution and geometry of karst features were observed. These include: (1) karst in Ordovician-age interbedded limestone-and-shale units is most developed south of the Illinoian glacial margin; (2) the number of sinkholes per square mile is greatest in areas where the bedrock is overlain by 50 to 100 feet of glacial drift; (3) karst development is most pronounced in the Columbus Limestone (Middle Devonian age) and Raccoon Creek Limestone (Middle Devonian) and Perdido Limestone (Lower Mississippian); and (4) sinkholes developed in susceptible carbonate units have surface expressions through as much as 20 feet of overlying noncarbonate materials such as shale or glacial drift.

FIGURE 3. Sinkholes develop through the following sequence of events: (1) Rain water reacts with carbon dioxide in the soil to form a weak carbonic acid ($H_2O + CO_2 \rightarrow H_2CO_3$). (2) Carbonic acid enters fractures in the bedrock and reacts with calcium carbonate ($H_2CO_3 + CaCO_3 \rightarrow Ca(CO_3)_2$), dissolving the rock and enlarging the fractures. (3) Water moves downward through the enlarged fractures, carrying with it the dissolved rock. (4) Over time, the fractures are enlarged to the point that the overlying soil can no longer bridge the gaps. (5) Soil collapses into the fractures, forming a sinkhole. Graphic from the Columbus Dispatch, used with permission.

FIGURE 4. With the passing of time, bedrock fractures become greatly enlarged by the dissolution and removal of carbonate rock. Eventually, the soil that fills the fractures is no longer capable of bridging the gaps and sink holes to collapse into the earth. Surface drainage is directed directly into the groundwater environment where solubility-induced disintegration, thereby accelerating the rate of sinkhole development. Eventually, the sinkhole will grow to the point that it will capture the entire stream and its surface flow through sinkhole. The stream vanishes and the presence of numerous sinkholes and other solution features is called karst. Graphic from the Columbus Dispatch, used with permission.

FIGURE 5. Over the course of many centuries, sinkholes continue to enlarge and coalesce with other sinkholes as underground water collapses and migrating streams and/or dissolution continues to remove rock. Eventually, the sinkhole will grow to the point that it will capture the entire stream and its surface flow into underground passage large enough for a person to enter. The entire lake continues to drain to sinkholes as chemical drainage networks widen the cave system. Sinkholes drain integrated and efficient karst. Over time, the cave ceilings and walls will continue to collapse, creating a valley in the overlying land. Cave ceilings and walls in streams along the cave floor may also form, causing water to be forced to the surface in springs. Graphic from the Columbus Dispatch, used with permission, modified slightly.

Drainage networks in karst terrains are commonly integrated and efficient. Over time, the cave ceilings and walls in streams along the cave floor may also form, causing water to be forced to the surface in springs. Graphic from the Columbus Dispatch, used with permission, modified slightly.

Probable karst areas were defined as areas that: (1) lie within a half mile of a known or indicated karst location, and (2) are underlain by carbonate or gypsiferous bedrock with overburden generally of less than 200 feet of non-carbonate bedrock and/or unconsolidated material as shown by comparison of 7.5-minute bedrock-topography and bedrock-geology maps to surface topography. Known karst locations and probable karst areas on the south terrain were digitized and are shown on this 1:500,000-scale map.

The May 2006 revision of this map is the first to portray information on the thickness of glacial drift and alluvium in areas of karst in Ohio. The drift-thickness map (figure 1, fig. 2) illustrates that karst terrains in Ohio is generally limited to areas of carbonate rock (limestone and dolomite) overlain by less than 200 feet of glacial drift and/or alluvium, and 2) isolated areas of the state where drift-thickness/bedrock-lithology conditions are most conducive to karst development, even though reconnaissance mapping may not have identified karst features. Areas where drift is thicker than 200 feet and areas where bedrock crops out are depicted on this map occur where bedrock is indicated to be carbonate by more than 200 feet of drift and/or by outside the boundaries of bedrock lithologies most susceptible to karst development. Such separate outliers from the bedrock used to map karst terrains are the Devonian age Ohio Shale and Silurian age Salina Group. Karst features were not available in 1997 when probability areas were originally determined, and 3) that have occasionally developed in areas where soluble bedrock underlies as much as 20 or more feet of relatively insoluble bedrock (e.g., noncalcareous shale).

Although Ohio has not been regarded as having a significant amount of karst, this investigation revealed many areas of the state that contain well-developed karst terrains, which occur mostly in four geographic regions of the state: (1) north-central Ohio, where a broad band of solution-collapse depressions extends from northeastern Seneca County to the Ohio River in Scioto County; (2) northeastern Ohio, in parts of Ashtabula, Lake, Geauga, Summit, and Portage counties; (3) southern Ohio, in carbonate outcrops lying beyond the Wisconsinan glacial boundary; and (4) west-central Ohio, in the Bellefontaine upland of Logan and Champaign Counties.

Several relationships between geologic units and the distribution and geometry of karst features were observed. These include: (1) karst in Ordovician-age interbedded limestone-and-shale units is most developed south of the Illinoian glacial margin; (2) the number of sinkholes per square mile is greatest in areas where the bedrock is overlain by 50 to 100 feet of glacial drift; (3) karst development is most pronounced in the Columbus Limestone (Middle Devonian age) and Raccoon Creek Limestone (Middle Devonian) and Perdido Limestone (Lower Mississippian); and (4) sinkholes developed in susceptible carbonate units have surface expressions through as much as 20 feet of overlying noncarbonate materials such as shale or glacial drift.

FIGURE 3. Sinkholes develop through the following sequence of events: (1) Rain water reacts with carbon dioxide in the soil to form a weak carbonic acid ($H_2O + CO_2 \rightarrow H_2CO_3$). (2) Carbonic acid enters fractures in the bedrock and reacts with calcium carbonate ($H_2CO_3 + CaCO_3 \rightarrow Ca(CO_3)_2$), dissolving the rock and enlarging the fractures. (3) Water moves downward through the enlarged fractures, carrying with it the dissolved rock. (4) Over time, the fractures are enlarged to the point that the overlying soil can no longer bridge the gaps. (5) Soil collapses into the fractures, forming a sinkhole. Graphic from the Columbus Dispatch, used with permission.

FIGURE 4. With the passing of time, bedrock fractures become greatly enlarged by the dissolution and removal of carbonate rock. Eventually, the soil that fills the fractures is no longer capable of bridging the gaps and sink holes to collapse into the earth. Surface drainage is directed directly into the groundwater environment where solubility-induced disintegration, thereby accelerating the rate of sinkhole development. Eventually, the sinkhole will grow to the point that it will capture the entire stream and its surface flow through sinkhole. The stream vanishes and the presence of numerous sinkholes and other solution features is called karst. Graphic from the Columbus Dispatch, used with permission.

FIGURE 5. Over the course of many centuries, sinkholes continue to enlarge and coalesce with other sinkholes as underground water collapses and migrating streams and/or dissolution continues to remove rock. Eventually, the sinkhole will grow to the point that it will capture the entire stream and its surface flow into underground passage large enough for a person to enter. The entire lake continues to drain to sinkholes as chemical drainage networks widen the cave system. Sinkholes drain integrated and efficient karst. Over time, the cave ceilings and walls will continue to collapse, creating a valley in the overlying land. Cave ceilings and walls in streams along the cave floor may also form, causing water to be forced to the surface in springs. Graphic from the Columbus Dispatch, used with permission, modified slightly.

KNOWN AND PROBABLE KARST IN OHIO DCMS 24 1999

Digital cartographic files of the locations of known and probable karst are merged with digital bedrock-geology maps to produce a non-GIS open-file map, DCMS 24, *Known and probable karst in Ohio*.



KNOWN AND PROBABLE KARST IN OHIO 2002

DCMS 24 is updated to true GIS format and renumbered EG-1